



**QUEEN'S  
UNIVERSITY  
BELFAST**

## **Appropriate treatment of depression in older community-dwelling adults**

Ryan, C. (2014). Appropriate treatment of depression in older community-dwelling adults. *Journal of Primary Health Care*, 6(4), 268. <https://www.rnzcgp.org.nz/assets/documents/Publications/JPHC/December-2014/JPHCGuestEditorialDecember2014.pdf>

**Published in:**  
Journal of Primary Health Care

**Document Version:**  
Publisher's PDF, also known as Version of record

**Queen's University Belfast - Research Portal:**  
[Link to publication record in Queen's University Belfast Research Portal](#)

### **General rights**

Copyright for the publications made accessible via the Queen's University Belfast Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

### **Take down policy**

The Research Portal is Queen's institutional repository that provides access to Queen's research output. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact [openaccess@qub.ac.uk](mailto:openaccess@qub.ac.uk).

# Appropriate treatment of depression in older community-dwelling adults

**Cristin Ryan** MPharm, PhD, MRPharmS

Lecturer in Clinical Pharmacy,  
School of Pharmacy, Queen's  
University Belfast, Belfast,  
United Kingdom

**T**he prevalence of depression increases with increasing age and is associated with a large disease burden and increased mortality.<sup>1</sup> Identifying, diagnosing and treating depression in older people (>65 years) is particularly important because the population is ageing progressively worldwide. However, the clinical management of depression in this age group is challenging, for several reasons. Firstly, older people have varied symptomatic presentations of depression, particularly if depression coexists with another age-related condition (e.g. Alzheimer's disease). Secondly, the selection of the most appropriate pharmacological treatment requires consideration of several factors, including the presence and treatment of coexisting morbidities, and age-related alterations in the pharmacokinetic and pharmacodynamic handling of medicines.<sup>2,3</sup>

As a result of these challenges, prescribing of medicines to treat depression in older people is not always entirely appropriate; treatment may be inappropriate, with both undertreatment and overtreatment problematic. Essentially, undertreatment can result from underdiagnosis and conservative prescribing, perhaps in an effort to keep a patient's drug burden to a minimum. Overtreatment on the other hand could be due to extended durations of treatment, or the prescribing of more medicines than actually indicated.<sup>4</sup>

There are several prescribing criteria developed to improve prescribing for older people, such as Beers' Criteria and STOPP/START.<sup>5,6</sup> Whilst these were developed as a guide to prescribing for older people in the general sense, they refer to instances where the prescribing of psychotropic medicines can be inappropriate. Indeed, the START<sup>6</sup> criteria also notes when prescribing of antidepressant medications should be considered.

Despite these criteria being readily available, several studies report the inappropriate use of psychotropic medicines in older people, particularly in those residing in nursing homes.

In order to improve prescribing for older, depressed, community-dwelling patients, the use of explicit prescribing criteria, as part of a medication review process, should be considered. Importantly, medication reviews facilitated by prescribing criteria should be conducted in tandem with diagnostic criteria and knowledge of patients' depressive symptoms. Further research is required to determine the long-term effects of inappropriate prescribing for depressive patients in primary care. Longitudinal cohort data would be ideal to meet this need.

## References

1. Rodda J, Walker Z, Carter J. Depression in older adults. *BMJ*. 2011;343:d5219.
2. Alexopoulos GS. Depression in the elderly. *Lancet*. 2005;365:1961–70.
3. Mangoni AA, Jackson SH. Age-related changes in pharmacokinetics and pharmacodynamics: basic principles and practical applications. *Br J Clin Pharmacol*. 2004;57(1):6–14.
4. O'Mahony D, Gallagher PF. Inappropriate prescribing in the older population: need for new criteria. *Age Ageing*. 2008;37:138–41.
5. Gallagher P, Ryan C, Byrne S, Kennedy J, O'Mahony D. STOPP (Screening Tool of Older Person's Prescriptions) and START (Screening Tool to Alert doctors to Right Treatment). Consensus validation. *Int J Clin Pharmacol Ther*. 2008;46:72–83.
6. American Geriatrics Society 2012 Beers Criteria Update Expert Panel. American Geriatrics Society updated Beers Criteria for potentially inappropriate medication use in older adults. *J Am Geriatr Soc*. 2012;60(4):616–631.

J PRIM HEALTH CARE  
2014;6(4):268.

## CORRESPONDENCE TO:

**Cristin Ryan**

School of Pharmacy,  
Queen's University Belfast,  
97 Lisburn Rd, Belfast  
BT73DE, United Kingdom  
c.ryan@qub.ac.uk